


U.S. Department of the Interior
U.S. Geological Survey

Beyond the Golden Gate— Oceanography, Geology, Biology, and Environmental Issues in the Gulf of the Farallones

Edited by
Herman A. Karl, John L. Chin, Edward Ueber,
Peter H. Stauffer, and James W. Hendley II

Circular 1198



U.S. Department of the Interior
Gale A. Norton, Secretary

U.S. Geological Survey
Charles G. Groat, Director

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.

U.S. Geological Survey, Reston, Virginia: 2001

For additional copies please contact:

USGS Information Services
Box 25286
Denver, CO 80225

This report and any updates to it are available at
<http://geopubs.wr.usgs.gov/circular/c1198/>

Additional USGS publications can be found at
<http://geology.usgs.gov/products.html>

For more information about the USGS and its products:

Telephone: 1-800-ASK-USGS (1-888-275-8747)
World Wide Web: <http://www.usgs.gov>

ISBN 0-607-95030-7

Published in the Western Region, Menlo Park, California
Manuscript approved for publication June 18, 2000

Production and design by Sara Boore and Susan Mayfield

Text and illustrations edited by Peter H. Stauffer,
James W. Hendley II, and George A. Havach

CD-ROM prepared by Michael F. Diggles

Cooperating Organizations

Bodega Marine Laboratory (University of California at Davis)

British Geological Survey

California Department of Health Services

Environmental Protection Agency

**Kernfysisch Versneller Instituut,
Groningen, The Netherlands**

**National Oceanic and Atmospheric
Administration**

*Gulf Of The Farallones National
Marine Sanctuary
Cordell Bank National Marine Sanctuary
National Marine Fisheries Service*

National Park Service

Point Reyes Bird Observatory

U.S. Army Corps of Engineers

U.S. Geological Survey

U.S. Navy

Contributors

Bodega Marine Laboratory (University of California at Davis)
Stephen R. Wing

British Geological Survey
David G. Jones
Philip D. Roberts

California Department of Health Services
Gregg W. Langlois
Patricia Smith

Environmental Protection Agency
Allan Ota

**Kernfysisch Versneller Instituut,
Groningen, The Netherlands**
Johannes Limburg

**National Oceanic and Atmospheric
Administration**

*Gulf of the Farallones National
Marine Sanctuary
Jan Roletto
Edward Ueber
Cordell Bank National Marine Sanctuary
Dan Howard
National Marine Fisheries Service
Peter Adams
Tom Laidig*

National Park Service
Scot Anderson

Point Reyes Bird Observatory
Scot Anderson
Peter Pyle

U.S. Geological Survey
John L. Chin
Walter E. Dean
James V. Gardner
Russell W. Graymer
James W. Hendley II
Herman A. Karl
Kaye Kinoshita
Marlene A. Noble
Stephanie L. Ross
Holly F. Ryan
William C. Schwab
Peter H. Stauffer
Florence L. Wong



Continental Slope Communities

Tom Laidig

In the shallow coastal areas of the Gulf of the Farallones, as in other regions of the world, fishing pressure has increased and numbers of fish have decreased over the past few decades. As many fish stocks have declined, some fishermen have been forced to look elsewhere to fill their nets.

Traditional fishing grounds in the gulf have been located on the Continental Shelf, a rather flat, relatively shallow area of the sea floor adjacent to the coast. At a depth of about 600 feet, the bottom starts to drop off more rapidly on what is called the Continental Slope. It is on upper and middle parts of this steeper slope that the new fishing grounds have been established. Because the fish inhabiting these deeper waters are less understood than those in shallower water, there is a danger of overharvesting, which could threaten the long-term viability of these newer fisheries.

The deep waters of the Continental Slope are characterized by nearly freezing temperatures, extremely low light conditions, and very high pressures. Because of the cold, organisms that live at these depths have slower metabolisms—they eat less frequently, are slower in digesting their food, and move and grow more slowly. They also attain greater ages than their counterparts that live in shallower

waters—some deep-sea rockfish live more than 70 years.

Many of the animals living in the perpetual darkness of the Continental Slope have developed light-producing organs. These serve various functions, such as communicating with members of their own kind (as in courtship), attracting food (like attracting moths to a flame), and avoiding being eaten (flashing a light in a predator's eyes can give an animal a chance to get away).

Another adaptation to the darkness is an absence of color diversity. With no light, colors have little function. Therefore, animals living on the Continental Slope are generally a dark color, like black, brown, or red. Among the fishes, rockfish and thornyheads are dominantly red. Red objects appear black at depth, allowing red organisms to blend in with their dark surroundings.

The water pressure on the sea floor at the top of the Continental Slope is more than 10 times higher than at the surface, and at the bottom of the slope the pressure can be more than 100 times higher than at the surface. To compensate for this high pressure, organisms have a large percentage of water in their tissues, bones, and shells that replaces other substances, such as gases and calcium. Owing to the high water content of

their tissues, many larger, older fish caught from deeper waters are limp and soft when brought to the surface.

Fishes living at different depths on the Continental Slope have different life histories. Species living near the top of the slope produce pelagic (open-ocean) young that spend the first few months to years of life swimming in the upper water column and then settle out in relatively shallow water and migrate downslope as they grow and mature. Dover sole, sablefish, and rockfish have this type of life history; however, most species living deeper, such as rattails, deep-sea soles, and slickheads, have young that live in the same depths as the adults.

Relatively few species occur at all or most depths on the Continental Slope. Species occupying one depth commonly are replaced by similar species at other depths. An exception is the eel-like hagfish, which is found at all depths on the slope.

Productive commercial fisheries operating today on the Continental Slope off California's coast catch Dover sole, sablefish, deep-living rockfishes, and thornyheads. Many of these fishes occupy similar habitats and generally are caught together. One increasingly active fishery is for rattails, a deep-living fish with a large head and a long tail that tapers to a point.

One major fishery of note is for hagfish, the skin of which is used to make what are sold as "eel skin" wallets. Hagfish are not true eels but are a primitive group of fish that have no bones or jaws. Instead of bones, they have cartilage, and instead of jaws, they have a large sucker-like mouth similar to that of a lamprey or a leech. Once attached, hagfish use a tongue with many tiny teeth to dig into their prey. Once inside, the prey is eaten from the inside out. Besides its unique method of eating, the hagfish has another interesting trait—it produces copious amounts of slime, probably used to discourage predators, which gives the fish its nickname, the "slime eel."

Except for fishing activities, the Continental Slope and its communities of fish and invertebrates are still virtually untouched by humans, offering scientists the opportunity to study a generally undisturbed natural system. New methods, such as viewing animals and their habitat by underwater video cameras in submarines and in remotely operated vehicles, have been particularly productive in providing a new understanding of fish and invertebrates living on the slope. In the Gulf of the Farallones, scientists are using these methods to collect data at increasingly greater depths, providing critical information needed to better protect these areas from overuse.




Northern lampfish with light organs (small white circles) along its underside. This small fish, only about 2.5 inches long, also displays the dark coloration typical of many animals on deeper parts of the Continental Slope. (Photograph courtesy of K. Sakuma and D. Roberts.)



Because of the absence of light on the Continental Slope, animals living there are generally either a dark color, such as the sablefish (left), or red, such as the shortspine thornyhead (right).






Juvenile rockfish swimming over Cordell Bank in the northern Gulf of the Farallones. Species of fish living near the top of the Continental Slope in the gulf produce pelagic (open-ocean) young that spend the first few months to years of life swimming in the upper water column and then settle out in relatively shallow water and migrate downslope as they grow and mature. (Photograph courtesy of Robert Schmieder, Cordell Expeditions.)



A red-banded rockfish on the Continental Slope in the Gulf of the Farallones. Relatively few species of fish occur at all or most depths on the slope. Those occupying one depth commonly are replaced by similar species at other depths. For example, greenstriped and stripetail rockfishes live on muddy bottoms on the upper part of the slope, whereas at greater depth they are replaced by species of thornyheads.



Thornyheads, giant brittle stars, and other animals on the deeper Continental Slope in the Gulf of the Farallones. (Photograph from Gulf of the Farallones National Marine Sanctuary.)